

On the cohomology of $sl(m+1, \mathbb{R})$ acting on differential operators and $sl(m+1, \mathbb{R})$ -equivariant symbol

P.B.A. Lecomte

Abstract

Let $\mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m)$ denote the space of differential operators of order $\leq k$ from the space of λ -densities of $\mathbb{R}^m$ into that of μ -densities and let $\mathcal{S}_{\delta}^k(\mathbb{R}^m)$ be the space of k -contravariant, symmetric tensor fields on $\mathbb{R}^m$ valued in the δ -densities, $\delta = \mu - \lambda$.

Denote by sl_{m+1} the projective embedding of $sl(m+1, \mathbb{R})$ as a subalgebra of the Lie algebra of vector fields of $\mathbb{R}^m$.

One computes the cohomology of sl_{m+1} with coefficients in the space of differential operators from $\mathcal{S}_{\delta}^p(\mathbb{R}^m)$ into $\mathcal{S}_{\delta}^q(\mathbb{R}^m)$. It is non vanishing only for some *critical* values of δ . For $m = 1$, these are the values pointed out by H. Gargoubi in a completely different context [6].

This allows to determine the condition under which the short exact sequence of sl_{m+1} -modules

$$0 \rightarrow \mathcal{D}_{\lambda\mu}^{k-1}(\mathbb{R}^m) \rightarrow \mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m) \xrightarrow{\sigma} \mathcal{S}_{\delta}^k(\mathbb{R}^m) \rightarrow 0$$

is split (σ is the symbol map).

From this one recovers and generalizes useful results about the structure of the sl_{m+1} -module $\mathcal{D}_{\lambda\mu}(\mathbb{R}^m) = \oplus_k \mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m)$ [3, 5, 6, 8].

The cohomology of the latter is also computed.

Key words : Lie algebra, differential operators, cohomology, infinitesimal homographies.

Classification numbers : 58B99, 17B56, 17B65, 47E05.

Titre courant : cohomology of $sl(m+1, \mathbb{R})$ on differential operators.

Rsum

Soient $\mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m)$ l'espace des opérateurs différentiels d'ordre $\leq k$ sur les λ -densités de $\mathbb{R}^m$ valeurs dans les μ -densités de $\mathbb{R}^m$ et $\mathcal{S}_{\delta}^k(\mathbb{R}^m)$ l'espace des champs de tenseurs k -contravariants symétriques sur $\mathbb{R}^m$ valeurs dans les δ -densités, o $\delta = \mu - \lambda$.

Soit galement sl_{m+1} le plongement projectif de $sl(m+1, \mathbb{R})$ dans l'algèbre de Lie des champs de vecteurs de $\mathbb{R}^m$.

On calcule la cohomologie de sl_{m+1} coefficients dans les opérateurs différentiels de $\mathcal{S}_{\delta}^p(\mathbb{R}^m)$ dans $\mathcal{S}_{\delta}^q(\mathbb{R}^m)$. Elle n'est non nulle que pour certaines valeurs,

dites *critiques*, de δ . Pour $m = 1$, ce sont les valeurs mises en vidence par H. Gargoubi dans un tout autre contexte [6].

A l'aide des rsultats obtenus, on dtermine quelle condition la courte suite exacte de sl_{m+1} -modules

$$0 \rightarrow \mathcal{D}_{\lambda\mu}^{k-1}(\mathbb{R}^m) \rightarrow \mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m) \xrightarrow{\sigma} \mathcal{S}_{\delta}^k(\mathbb{R}^m) \rightarrow 0$$

est scinde (σ est le passage au symbole).

Cela permet de retrouver et de gnraliser diffrents rsultats utiles concernant la structure de sl_{m+1} -module de $\mathcal{D}_{\lambda\mu}(\mathbb{R}^m) = \bigoplus_k \mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m)$ [3, 5, 6, 8].

La cohomologie de ce dernier est galement calcule.

1 Introduction

Let $\mathcal{D}_{\lambda\mu}(M)$ denote the space of differential operators from the space of scalar λ -densities over a smooth manifold M into the space of scalar μ -densities.

The Lie derivation with respect to vector fields equips $\mathcal{D}_{\lambda\mu}(M)$ with a natural structure of module over the Lie algebra of vector fields $Vect(M)$ of M . The study of that module has been started in [3, 5, 6, 8, 9]. See also [2] for pseudodifferential operators on the complex line. As a vector space, $\mathcal{D}_{\lambda\mu}(M)$ is isomorphic to the graded space

$$\mathcal{S}_{\delta}(M) = \bigoplus_{k \geq 1} \frac{\mathcal{D}_{\lambda\mu}^k(M)}{\mathcal{D}_{\lambda\mu}^{k-1}(M)}$$

associated to the filtration of $\mathcal{D}_{\lambda\mu}(M)$ by the order k of differentiation. The k -th term of this sum is nothing else but the space $\mathcal{S}_{\delta}^k(M) = \Gamma(\vee^k TM \otimes \Delta^{\delta} M)$ of k -symmetric contravariant tensor fields valued in the scalar densities of weight $\delta = \mu - \lambda$ ($\Delta^{\delta} M$ denotes the vector bundle of δ -densities of M). The isomorphism between $\mathcal{D}_{\lambda\mu}^k(M)/\mathcal{D}_{\lambda\mu}^{k-1}(M)$ and $\mathcal{S}_{\delta}^k$ is of course induced in a natural way by the symbol map

$$\sigma : \mathcal{D}_{\lambda\mu}^k(M) \rightarrow \mathcal{S}_{\delta}^k(M).$$

This map commutes with the Lie derivation of operators on the left and of tensor fields on the right. But, however, $\mathcal{D}_{\lambda\mu}(M)$ and $\mathcal{S}_{\delta}(M)$ are not isomorphic as $Vect(M)$ -modules : $\mathcal{D}_{\lambda\mu}(M)$ is better viewed as a non trivial deformation of $\mathcal{S}_{\delta}(M)$.

For $\lambda = \mu$ and $m > 1$, it has been shown [8], nevertheless, that $\mathcal{D}_{\lambda\mu}(M)(\mathbb{R}^m)$ and $\mathcal{S}_0(\mathbb{R}^m)$ are isomorphic as sl_{m+1} -modules, where $sl_{m+1} \subset Vect(\mathbb{R}^m)$ is the canonical embedding of $sl(m+1, \mathbb{R})$ as the algebra of infinitesimal linear fractional local transformations of $\mathbb{R}^m$ (note that it is a maximal subalgebra of the algebra of polynomial vector fields on $\mathbb{R}^m$).

A similar result has been obtained in the one dimensional case (cf. [2] for pseudodifferential operators, [6] for differential operators) : except for special critical values of λ and μ , $\mathcal{D}_{\lambda\mu}(M)$ is again sl_2 -isomorphic to $\mathcal{S}_{\delta}(M)$. For the critical values, $\mathcal{D}_{\lambda\mu}(M)$ is a non trivial deformation of $\mathcal{S}_{\delta}(M)$ and it is described in [6] in terms of a family of cocycles, taylored on the occasion, of sl_2 acting on the space of differential operators from $\Gamma(\Delta^{-n/2} M)$ into $\Gamma(\Delta^{(n+2)/2} M)$.

In both cases, $m = 1$ and $m > 1$, the sl_{m+1} -module structure of $\mathcal{D}_{\lambda\mu}(\mathbb{R}^m)$ proved to be a powerful tool in studying its $Vect(\mathbb{R}^m)$ -structure.

In some sense, the possible difference between the L -modules $\mathcal{D}_{\lambda\mu}(M)$ and $\mathcal{S}_\delta(M)$, $L = Vect(M)$ or $s\ell_{m+1}$, follows from the fact that the short exact sequence of modules

$$0 \rightarrow \mathcal{D}_{\lambda\mu}^{k-1}(M) \xrightarrow{i} \mathcal{D}_{\lambda\mu}^k(M) \xrightarrow{\sigma} \mathcal{S}_\delta^k(M) \rightarrow 0 \quad (1)$$

(i : the inclusion) maybe is not split. The obstruction against splitting is a member of

$$H^1(L, Hom(\mathcal{S}_\delta^k(M), \mathcal{D}_{\lambda\mu}^{k-1}(M))).$$

Composing with the symbol map $\sigma : \mathcal{D}_{\lambda\mu}^{k-1}(M) \rightarrow \mathcal{S}_\delta^{k-1}(M)$, one gets a sort of first order approximation of this obstruction, namely an element of

$$H^1(L, Hom(\mathcal{S}_\delta^k(M), \mathcal{S}_\delta^{k-1}(M))).$$

In this paper, we compute the cohomology space

$$H(s\ell_{m+1}, \mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))).$$

This allows us not only to recover the critical values of [6] but also to find critical values for $m > 1$. To do that, we compute the spaces

$$H(s\ell_{m+1}, \mathcal{D}^k(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)))$$

($\mathcal{D}^k(A, B)$ denotes the space of differential operators of order $\leq k$ from A into B).

They are computed by induction on k , using the short exact sequence associated to the inclusion of the operators of order $k-1$ into these of order k . The critical values occur when $p-q \in \mathbb{N}_0$. They are then these for which the connecting homomorphism of the corresponding exact sequence in cohomology vanishes. Equivalently, they are these for which

$$H(s\ell_{m+1}, \mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))) \neq 0.$$

We also compute the space $Hom_{s\ell_{m+1}}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$. This is quite hard because one is left to show that a mapping from $\mathcal{S}_\delta^p(\mathbb{R}^m)$ into $\mathcal{S}_\delta^q(\mathbb{R}^m)$ is local knowing only that it is commuting with the L_X , $X \in s\ell_{m+1}$. As a corollary, we get generalization of a result of [8] : if $(m+1)\delta - m \notin \mathbb{N}_0$, then a linear map from $\mathcal{S}_\delta(\mathbb{R}^m)$ is $s\ell_{m+1}$ -equivariant if and only if its restriction to each $\mathcal{S}_\delta^k(\mathbb{R}^m)$, $k \in \mathbb{N}$, is a constant multiple of the identity.

As a consequence of our computations, we find the necessary and sufficient condition for (1) to be split as a sequence of $s\ell_{m+1}$ -modules (with $M = \mathbb{R}^m$). In particular, we show that if $(m+1)\delta - m \notin \mathbb{N}_0$, then the $s\ell_{m+1}$ -modules $\mathcal{D}_{\lambda\mu}(\mathbb{R}^m)$ and $\mathcal{S}_\delta(\mathbb{R}^m)$ are canonically isomorphic. This also generalizes a useful result of [8].

In fact, the above mentioned cohomology classes constructed in [6] are valued in $\mathcal{D}_{-\frac{n}{2}, 1+\frac{n}{2}}(\mathbb{R})$. We recover them as a consequence of our computations due to the fact that $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}), \mathcal{S}_\delta^q(\mathbb{R}))$ is isomorphic to $\mathcal{D}_{\delta-p, \delta-q}(\mathbb{R}^m)$.

Besides, very little more is needed then to compute the cohomology of $s\ell_{m+1}$ with coefficients in $\mathcal{D}_{\lambda\mu}(\mathbb{R}^m)$, $m \geq 1$. This is done in the last section. The difference between the cases $m > 1$ and $m = 1$ is then really significant.

2 The projective embedding of $sl(m+1, \mathbb{R})$

For each $T \in sl(m+1, \mathbb{R})$ we denote by T^* the fundamental vector fields on $P_m\mathbb{R}$ associated to T and to the canonical action of $SL(m+1, \mathbb{R})$ by linear projective transformations. We also denote by T^* its restriction to $\mathbb{R}^m$, viewed as the open set of points of $P_m\mathbb{R}$ having a non vanishing $(m+1)$ -th projective coordinate.

The mapping $T \rightarrow T^*$ is an injective homomorphism of Lie algebras from $sl(m+1, \mathbb{R})$ into the Lie algebra $Vect(\mathbb{R}^m)$ of vector-fields on $\mathbb{R}^m$.

Following [8], we call it the *projective embedding of $sl(m+1, \mathbb{R})$* . We now describe it in a way well suited to our purpose.

We realize $sl(m+1, \mathbb{R})$ as $\mathbb{R}^m \oplus gl(m, \mathbb{R}) \oplus \mathbb{R}^{m*}$, where $\mathbb{R}^m$ and $\mathbb{R}^{m*}$ are abelian subalgebras, where the adjoint action of $A \in gl(m, \mathbb{R})$ is the natural representation of $gl(m, \mathbb{R})$ and where the bracket of $h \in \mathbb{R}^m$ and $\alpha \in \mathbb{R}^{m*}$ is given by

$$[h, \alpha] = \alpha(h)\mathbf{1} + h \otimes \alpha$$

($\mathbf{1}$ is the unit matrix). It is easily seen that the fundamental vector fields associated to $h = (h^i) \in \mathbb{R}^m$, $A = (A_j^i) \in gl(m, \mathbb{R})$ and $\alpha = (\alpha_i) \in \mathbb{R}^{m*}$ are

$$h^* = -h^i \partial_i, \quad A^* = -A_j^i x^j \partial_i \quad \text{and} \quad \alpha^* = \alpha(x) x^i \partial_i$$

(sums over repeated indices are understood and ∂_i denotes the partial derivative with respect to x^i).

Note that $sl(m+1, \mathbb{R})$ is a graded Lie algebra, $\mathbb{R}^m$, $gl(m, \mathbb{R})$ and $\mathbb{R}^{m*}$ being the homogeneous components of degree $-1, 0$ and 1 respectively.

For the sake of brevity, we denote sl_{m+1} the above realization of $sl(m+1, \mathbb{R})$.

3 Cohomology of sl_{m+1} associated to a representation of $gl(m, \mathbb{R})$

Let (V, ρ) be a finite dimensional representation of $gl(m, \mathbb{R})$.

The space $C_\infty(\mathbb{R}^m, V)$ of smooth V -valued functions on $\mathbb{R}^m$ is a representation of $Vect(\mathbb{R}^m)$. The action L_X^ρ of a vector field X on a function f is given by

$$L_X^\rho f = X.f - \rho(DX)f$$

where $X.f$ is the usual derivative $X^i \partial_i f$ and where $DX = (\partial_j X^i)$ is the differential of X .

In fact, $C_\infty(\mathbb{R}^m, V)$ is the space of sections of the (trivial) bundle associated to the linear frame bundle of $\mathbb{R}^m$ and to V and L^ρ is the corresponding natural Lie derivation.

Our purpose is to compute the Chevalley-Eilenberg cohomology of the restriction of L^ρ to sl_{m+1} .

Recall that L^ρ induces a representation of $\mathcal{L}^\rho$ on the space of cochains. It is defined by

$$\mathcal{L}_X^\rho = i_X \circ \partial^\rho + \partial^\rho \circ i_X, \quad \forall X \in sl_{m+1},$$

where ∂^ρ is the coboundary operator. One has also

$$(\mathcal{L}_X^\rho c)(X_0, \dots, X_{s-1}) = L_X^\rho(c(X_0, \dots, X_{s-1})) - \sum_{0 \leq i < s} c(X_0, \dots, [X, X_i], \dots, X_{s-1})$$

for each s -cochain c .

Proposition 3.1 *For each $u \in \{0, \dots, m\}$, there exists a unique linear $\chi^u : \Lambda(\mathfrak{gl}(m, \mathbb{R}), \Lambda^u(\mathbb{R}^{m*}, V)) \rightarrow \Lambda(\mathfrak{sl}_{m+1}, C_\infty(\mathbb{R}^m, V))$ such that*

- (i) $\mathcal{L}_{h^*}^\rho \circ \chi^u = 0$, $i_{h^*} \circ \chi^u = 0$, $\forall h \in \mathbb{R}^m$,
- (ii) $(\chi^u(\gamma))(A_0^*, \dots, A_{t-1}^*, \alpha_1^*, \dots, \alpha_u^*) = (\gamma(A_0, \dots, A_{t-1}))(\alpha_1, \dots, \alpha_u)$ for all $A_i \in \mathfrak{gl}(m, \mathbb{R})$ and all $\alpha_j \in \mathbb{R}^{m*}$,
- (iii) $(\chi^u(\gamma))(X_0, \dots, X_{t+u-1})$ is an homogeneous polynomial of degree $-u + \sum_i \text{degree}(X_i)$, for all $X_i \in \mathfrak{sl}_{m+1}$.

Proof. It is easy to verify that the mapping that sends γ onto the cochain

$$(X_0, \dots, X_{t+u-1}) \rightarrow \frac{(-1)^t}{t!u!(m+1)^u} \sum_{\nu} \text{sign}(\nu) (\gamma(DX_{\nu_0}, \dots, DX_{\nu_{t-1}}))(d \text{tr}(DX_{\nu_t}), \dots, d \text{tr}(DX_{\nu_{t+u-1}}))$$

has the required properties (ν describes the permutations of $t+u$ elements and $\text{sign}(\nu)$ is the signature of the permutation ν). Hence the existence.

Let $c \in \Lambda^{t+u}(\mathfrak{sl}_{m+1}, C_\infty(\mathbb{R}^m, V))$ be such that $\mathcal{L}_{h^*} c = 0$, $i_{h^*} c = 0$ for each $h \in \mathbb{R}^m$. Suppose also that $c(X_0, \dots, X_{t+u-1})$ is an homogeneous polynomial of degree $-u + \sum_i \text{degree}(X_i)$ for all X_i . We claim that if, in addition,

$$c(A_0^*, \dots, A_{t-1}^*, \alpha_1^*, \dots, \alpha_u^*) = 0$$

for all $A_i \in \mathfrak{gl}(m, \mathbb{R})$ and all $\alpha_j \in \mathbb{R}^{m*}$, then $c = 0$, thus proving the uniqueness of χ^u . We show that $c(X_0, \dots, X_{t+u-1}) = 0$ by induction on $k = -u + \sum_i \text{degree}(X_i)$. It is true for $k \leq 0$, by assumption. If it is true for $k < \ell$ then for $k = \ell$, it follows from $\mathcal{L}^\rho c = 0$ that

$$h^*.(c(X_0, \dots, X_{t+u-1})) = 0, \quad \forall h \in \mathbb{R}^m.$$

The homogeneous polynomial $c(X_0, \dots, X_{t+u-1})$ of degree $\ell > 0$ is thus constant. Hence it vanishes. $\blacksquare$

Let χ denote the mapping from $\Lambda(\mathfrak{gl}(m, \mathbb{R}), \Lambda(\mathbb{R}^{m*}, V))$ into $\Lambda(\mathfrak{sl}_{m+1}, C_\infty(\mathbb{R}^m, V))$ that reduces to χ^u when restricted to $\Lambda(\mathfrak{gl}(m, \mathbb{R}), \Lambda^u(\mathbb{R}^m, V))$. Observe also that $\Lambda(\mathbb{R}^{m*}, V)$ is a representation of $\mathfrak{gl}(m, \mathbb{R})$, its action being deduced in a natural way from the given ρ and the natural action on $\mathbb{R}^{m*}$. We denote also by ∂^ρ the Chevalley-Eilenberg coboundary operator associated to that representation.

Lemma 3.2 *The mapping χ is an homomorphism of differential spaces.*

Proof. Let γ be given. Then $\mathcal{L}_{h^*}^\rho \partial^\rho \chi^u(\gamma) = \partial^\rho \mathcal{L}_{h^*}^\rho \chi^u(\gamma) = 0$ and $i_{h^*} \partial^\rho \chi^u(\gamma) = \mathcal{L}_{h^*}^\rho \chi^u(\gamma) - \partial^\rho i_{h^*} \chi^u(\gamma) = 0$. In addition, it is clear that

$$(\partial^\rho \chi^u(\gamma))(A_0^*, \dots, A_t^*, \alpha_1^*, \dots, \alpha_u^*) = ((\partial^\rho \gamma)(A_0, \dots, A_t))(\alpha_1, \dots, \alpha_u)$$

for all $A_i \in g\ell(m, \mathbb{R})$ and all $\alpha_j \in \mathbb{R}^{m^*}$. Finally, a careful examination of the various terms of $(\partial^\rho \chi^u(\gamma))(X_0, \dots, X_{t+u})$ shows that it is an homogeneous polynomial of degree $-u + \sum_i \text{degree}(X_i)$. In view of the uniqueness property of χ^u , it follows that $\partial^\rho \chi^u(\gamma) = \chi^u(\partial^\rho \gamma)$. ■

We are now in position to compute the cohomology of $s\ell_{m+1}$ associated to the representation L^ρ .

Theorem 3.3 *The mapping*

$$\chi_\sharp : H(g\ell(m, \mathbb{R}), \Lambda(\mathbb{R}^{m^*}, V)) \rightarrow H(s\ell_{m+1}, C_\infty(\mathbb{R}^m, V))$$

is a bijection.

Proof. We use the Hochschild-Serre spectral sequence $E_i^{p,q}$ associated to the subalgebra $\mathbb{R}^m$ of $s\ell_{m+1}$. Recall [4] that $E_1^{p,q}$ is the q -th cohomology space of the Chevalley-Eilenberg complex of the representation of $\mathbb{R}^m$ induced by $h \rightarrow \mathcal{L}_{h^*}^\rho$ on the space $\Lambda^p(s\ell_{m+1}/\mathbb{R}^m, C_\infty(\mathbb{R}^m, V))$ or, equivalently, on the space of the elements of $\Lambda^p(s\ell_{m+1}, C_\infty(\mathbb{R}^m, V))$ that are cancelled out by each i_{h^*} , $h \in \mathbb{R}^m$.

Once evaluated on $\mathbf{X} = (X_1, \dots, X_p) \in s\ell_{m+1}^p$, a q -cochain c defines a differential q -form on $\mathbb{R}^m$, namely

$$\omega_{\mathbf{X}}^c : (h_1, \dots, h_q) \rightarrow (c(h_1^*, \dots, h_q^*))(X_1, \dots, X_p).$$

As easily seen, if $\omega_{\mathbf{X}}^c = 0$ for degree $(\mathbf{X}) = \sum_i \text{degree}(X_i) < u$, then for degree $(\mathbf{X}) = u$, one has

$$d\omega_{\mathbf{X}}^c = \omega_{\mathbf{X}}^{\partial^\rho c},$$

where d is the de Rham differential. An easy induction on degree (X) allows thus to show that

(a) *If $q > 0$, then $E_1^{p,q} = 0$.*

Now, a 0-cocycle is an element $c \in \Lambda^p(s\ell_{m+1}, C_\infty(\mathbb{R}^m, V))$ such that $\mathcal{L}_{h^*}^\rho c = 0$ and $i_{h^*} c = 0$ for each $h \in \mathbb{R}^m$. In particular, if $c(\mathbf{X}) = 0$ when degree $(\mathbf{X}) < u$, then $c(\mathbf{X})$ is constant for degree $(\mathbf{X}) = u$. Thus, if $\gamma \in \Lambda^{p-u}(g\ell(m, \mathbb{R}), \Lambda^u(\mathbb{R}^{m^*}, V))$ is defined by

$$(\gamma(A_1, \dots, A_{p-u}))(\alpha_1, \dots, \alpha_u) = c(A_1^*, \dots, A_{p-u}^*, \alpha_1^*, \dots, \alpha_u^*)$$

then $c - \chi^u(\gamma)$ vanishes on arguments $\mathbf{X}$ of degree $\leq u$. It follows by induction on u that c belongs to the image of χ . It is clear that χ is injective. Therefore,

(b) $\chi : \Lambda(g\ell(m, \mathbb{R}), \Lambda(\mathbb{R}^{m^*}, V)) \rightarrow \oplus_{p \geq 0} E_1^{p,0}$ *is an isomorphism of differential spaces.*

We deduce from (a) that the higher differentials d_i , $i > 1$, of the spectral sequence are vanishing. The result then follows from (b). ■

4 Deforming the representations of $gl(m, \mathbb{R})$

Let again (V, ρ) be a finite dimensional representation of $gl(m, \mathbb{R})$. For each $\lambda \in \mathbb{R}$, define ρ_λ by

$$\rho_\lambda : A \rightarrow \rho(A) - \lambda \operatorname{tr}(A)id$$

where tr denotes the trace and id the identity from V into V .

We want to compute the cohomology of the representation (V, ρ_λ) of $gl(m, \mathbb{R})$ in terms of that of (V, ρ) .

Let $\Phi : (\mathcal{A}, \partial) \rightarrow (\mathcal{B}, \partial)$ be an homomorphism of differential spaces. The mapping

$$\partial_\Phi : (a, b) \mapsto (\partial a, \Phi(a) - \partial b)$$

is a differential on $\mathcal{A} \oplus \mathcal{B}$.

Lemma 4.1 *The cohomology space $H(\mathcal{A} \oplus \mathcal{B}, \partial_\Phi)$ is isomorphic to $\ker \Phi_\# \oplus H(\mathcal{B}, \partial)/\operatorname{im} \Phi_\#$.*

Proof. Indeed, the connecting homomorphism ([7]) of the short exact sequence of differential spaces

$$0 \rightarrow \mathcal{B} \xrightarrow{i} \mathcal{A} \oplus \mathcal{B} \xrightarrow{j} \mathcal{A} \rightarrow 0$$

where $iy = (0, -y)$ and $j(x, y) = x$, is $\Phi_\#$. ■

Proposition 4.2 *The Chevalley-Eilenberg cohomology of (V, ρ_λ) is isomorphic to*

$$\ker \rho_\lambda(\mathbf{1})_\# \oplus H(sl(m, \mathbb{R}), V)/\operatorname{im} \rho_\lambda(\mathbf{1})_\#$$

where $\rho_\lambda(\mathbf{1})_\#$ is the map induced in cohomology by

$$c \in \Lambda(sl(m, \mathbb{R}), V) \rightarrow \rho_\lambda(\mathbf{1}) \circ c \in \Lambda(sl(m, \mathbb{R}), V).$$

Proof. Indeed, the map

$$c \rightarrow (c|_{sl(m, \mathbb{R})}, (i_1 c)|_{sl(m, \mathbb{R})}) \tag{2}$$

is an isomorphism of differential spaces between $\Lambda(gl(m, \mathbb{R}), V)$ equipped with the differential ∂^{ρ_λ} and $(\Lambda sl(m, \mathbb{R}))^2$ equipped with the differential

$$(c', c'') \rightarrow (\partial^\rho c', \rho_\lambda(\mathbf{1}) \circ c' - \partial^\rho c'').$$

Hence the result, in view of Lemma 4.1. ■

We shall now apply the above proposition when $V \subset \otimes_b^a \mathbb{R}^m$ and when ρ is the natural representation of $gl(m, \mathbb{R})$ on the tensors. For the sake of simplicity, V_λ will denote the representation (V, ρ_λ) .

Proposition 4.3 *Let V be a subspace of $\otimes_b^a \mathbb{R}^m$ stable under the natural representation of $gl(m, \mathbb{R})$ on the tensors. If $\lambda \neq (a - b)/m$, then $H(gl(m, \mathbb{R}), V_\lambda) = 0$. If $\lambda = (a - b)/m$, then $H(gl(m, \mathbb{R}), V_\lambda)$ is isomorphic to*

$$(\Lambda gl(m, \mathbb{R})^*)_{g-inv} \otimes V_{s-inv} \tag{3}$$

where $g-inv$ and $s-inv$ denote the invariance with respect to $gl(m, \mathbb{R})$ and $sl(m, \mathbb{R})$ respectively.

Proof. If $m > 1$, the space of cocycles of the Chevalley-Eilenberg cohomology of the representation (V, ρ) of $sl(m, \mathbb{R})$ is the direct sum of the space of coboundaries and of $(\Lambda sl(m, \mathbb{R})^*)_{s-inv} \otimes V_{s-inv}$ [4]. On the latter, $\rho_\lambda(\mathbf{1})_\#$ is the multiplication by $a - b - \lambda m$. Moreover, (2) pulls $(\Lambda sl(m, \mathbb{R})^*)_{s-inv}^2$ back onto $(\Lambda gl(m, \mathbb{R})^*)_{g-inv}$. Hence the result when $m > 1$. For $m = 1$, the result follows from an immediate direct computation, (3) being just V in this case. ■

Corollary 4.4 *Under the assumptions of Proposition 4.3, $H(gl(m, \mathbb{R}), V_\lambda) \neq 0$ only if $\lambda = (a - b)/m$ is an integer.*

Proof. It is obvious if $m = 1$. If $m > 1$, it follows from the description of $(\otimes_b^a \mathbb{R}^m)_{s-inv}$ [11] : $T \in \otimes_b^a \mathbb{R}^m$ is $sl(m, \mathbb{R})$ -invariant if and only if $T(\xi_1, \dots, \xi_a, X_1, \dots, X_b)$, as a function of $\xi_i \in \mathbb{R}^{m*}$ and $X_j \in \mathbb{R}^m$, is a linear combination of products of contractions $\langle X_j, \xi_i \rangle$ and of determinants of the form $\det(\xi_{i_1}, \dots, \xi_{i_m})$ and $\det(X_{j_1}, \dots, X_{j_m})$. ■

5 Two examples

In this section, we illustrate the previous results by computing two cohomologies of sl_{m+1} that will be needed later.

Let us first consider $S_\lambda^k \mathbb{R}^m$, the space $\vee^k \mathbb{R}^m$ of k -contravariant symmetric tensors equipped with the representation ρ_λ (where ρ is the natural representation of $gl(m, \mathbb{R})$ on tensors).

Proposition 5.1 (i) *Assume that $m > 1$. If $(\lambda, k) \in \{(0, 0), (1, 0), (1, 1)\}$, then $H(sl_{m+1}, C_\infty(\mathbb{R}^m, S_\lambda^k \mathbb{R}^m))$ is isomorphic to $(\Lambda gl(m, \mathbb{R})^*)_{g-inv}$ otherwise it is vanishing.*

(ii) *If $(\lambda, u) \in \{(k, 0), (k, 1), (k + 1, 1), (k + 1, 2)\}$ then $H(sl_2, C_\infty(\mathbb{R}, S_\lambda^k \mathbb{R}))$ is isomorphic to $\mathbb{R}$ otherwise it vanishes.*

Proof. It follows from Theorem 3.3 and Proposition 4.3 that $H^u(sl_{m+1}, C_\infty(\mathbb{R}^m, S_\lambda^k \mathbb{R}^m))$ is non vanishing only if $\lambda = (j + k)/m$ for some $j \in \{0, \dots, m\}$ and that it is then isomorphic to

$$(\Lambda^{u-j} gl(m, \mathbb{R})^*)_{g-inv} \otimes \Lambda^j(\mathbb{R}^{m*}, \vee^k \mathbb{R}^m)_{s-inv}. \quad (4)$$

Using the description of $(\otimes_b^a \mathbb{R}^m)_{s-inv}$ recalled in the proof of Corollary 4.4, the latter is non vanishing if and only if (j, k) equals $(0, 0)$, $(m, 0)$ or $(m - 1, 1)$ in which case, (4) is respectively spanned by the mappings

$$\begin{aligned} (A_0, \dots, A_{t-1}) &\rightarrow \gamma(A_0, \dots, A_{t-1}), \\ (A_0, \dots, A_{t-1}, \alpha_1, \dots, \alpha_m) &\rightarrow \gamma(A_0, \dots, A_{t-1}) \det(\alpha_1, \dots, \alpha_m) \end{aligned}$$

or

$$(A_0, \dots, A_{t-1}, \alpha_1, \dots, \alpha_{m-1}) \rightarrow [\xi \rightarrow \gamma(A_0, \dots, A_{t-1}) \det(\alpha_1, \dots, \alpha_{m-1}, \xi)]$$

($t = u - j$, $A_i \in gl(m, \mathbb{R})$, $\alpha_i, \xi \in \mathbb{R}^{m*}$, $\gamma \in (\Lambda^t gl(m, \mathbb{R})^*)_{g-inv}$). Hence (i).

For (ii), it suffices to note that $\Lambda^j(\mathbb{R}, \vee^t \mathbb{R})_{s-inv} = \mathbb{R}$. ■

Corollary 5.2 *The cohomology of sl_{m+1} acting on the space of scalar λ -densities of $\mathbb{R}^m$ vanishes if $\lambda \neq 0$ and $\lambda \neq 1$ otherwise, it is isomorphic to $(\Lambda gl(m, \mathbb{R})^*)_{g-inv}$.*

Proof. It is the cohomology of the representation $C_\infty(\mathbb{R}^m, S_\lambda^0 \mathbb{R}^m)$ of sl_{m+1} . ■

Let us now consider the case of the space $S_p^{k,q} \mathbb{R}^m$ of k -contravariant symmetric tensors valued in the space of linear maps from $\vee^p \mathbb{R}^m$ into $\vee^q \mathbb{R}^m$:

$$\begin{aligned} S_p^{k,q} \mathbb{R}^m &= \vee^k \mathbb{R}^m \otimes Hom(\vee^p \mathbb{R}^m, \vee^q \mathbb{R}^m) \\ &\simeq \vee^k \mathbb{R}^m \otimes \vee^q \mathbb{R}^m \otimes \vee^p \mathbb{R}^{m*}. \end{aligned}$$

We equip it with the natural representation of $gl(m, \mathbb{R})$.

Proposition 5.3 *If $p \in \{k+q, k+q+1\}$ then $H(sl_{m+1}, C_\infty(\mathbb{R}^m, S_p^{k,q} \mathbb{R}^m))$ is isomorphic to $(\Lambda gl(m, \mathbb{R})^*)_{g-inv}$ otherwise it vanishes.*

Proof. The proof is similar to that of the previous proposition. One is now left to study

$$(\Lambda^{u-j} gl(m, \mathbb{R})^*)_{g-inv} \otimes \Lambda^j(\mathbb{R}^{m*}, S_p^{k,q} \mathbb{R}^m)_{s-inv} \quad (5)$$

knowing that $p = j + k + q$ (since “ λ ” = 0). This space is non vanishing if and only if $j = 0$ or $j = 1$, (5) being then respectively spanned by the mappings

$$(A_0, \dots, A_{t-1}) \rightarrow [P \rightarrow \gamma(A_0, \dots, A_{t-1})(\eta D_\xi)^k P]$$

or

$$(A_0, \dots, A_{t-1}, \alpha) \rightarrow [P \rightarrow \gamma(A_0, \dots, A_{t-1})(\alpha D_\xi)(\eta D_\xi)^k P]$$

($t = u - j$, $A_i \in gl(m, \mathbb{R})$, $\alpha \in \mathbb{R}^m$, $\gamma \in (\Lambda^t gl(m, \mathbb{R})^*)_{g-inv}$, $P \in \vee^k \mathbb{R}^m$).

Here, we view an element of $S_p^{k,q} \mathbb{R}^m$ as being a homogeneous polynomial of degree k in $\eta \in \mathbb{R}^{m*}$ valued in the space of linear mappings from the space of homogeneous polynomials of degree p in $\xi \in \mathbb{R}^{m*}$ into the space of these of degree q . Moreover, αD_ξ and ηD_ξ denote the derivations with respect to ξ in the direction $\alpha \in \mathbb{R}^{m*}$ and $\eta \in \mathbb{R}^{m*}$ respectively. These conventions will be helpfull in the sequel. ■

6 Cohomology of sl_{m+1} valued in $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$

Recall from the introduction that the structure of the sl_{m+1} -module $\mathcal{D}_{\lambda\mu}(\mathbb{R}^m)$ is related to the cohomology of sl_{m+1} with coefficients in $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$, $\delta = \mu - \lambda$.

We first compute the cohomology of sl_{m+1} with coefficients in $\mathcal{D}^k(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$, $k \in \mathbb{N}$. We proceed by induction on k , using the short exact sequence

$$\begin{aligned} 0 \rightarrow \mathcal{D}^{k-1}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)) &\xrightarrow{i} \mathcal{D}^k(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)) \\ &\xrightarrow{\sigma} C_\infty(\mathbb{R}^m, S_p^{k,q} \mathbb{R}^m) \rightarrow 0 \end{aligned}$$

that induces an exact triangle

$$\begin{array}{ccc}
H(s\ell_{m+1}, \mathcal{D}^{k-1}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))) & & \\
\theta \uparrow & \searrow i_\# & \\
H(s\ell_{m+1}, C_\infty(\mathbb{R}^m, S_p^{k,q}\mathbb{R}^m)) & \swarrow \sigma_\# & H(s\ell_{m+1}, \mathcal{D}^k(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))) \quad (6)
\end{array}$$

where the connecting homomorphism θ is of degree 1 ([7]). Note that the cohomology of the quotient is that computed in the previous section. From Proposition 5.3, we immediately deduce

Theorem 6.1 (a) If $p < q$, then $H(s\ell_{m+1}, \mathcal{D}^k(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))) = 0$
(b) If $n \geq 0$, then
(i) $k < n - 1 \Rightarrow H(s\ell_{m+1}, \mathcal{D}^k(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))) = 0$
(ii) $H(s\ell_{m+1}, \mathcal{D}^{n-1}(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)))$ is isomorphic to $(\Lambda gl(m, \mathbb{R})^*)_{g-inv}$
(iii) The inclusion of $\mathcal{D}^n(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ into $\mathcal{D}^k(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ induces an isomorphism

$$H(s\ell_{m+1}, \mathcal{D}^n(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))) \rightarrow H(s\ell_{m+1}, \mathcal{D}^k(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)))$$

for each $k > n$.

Remark 6.2 In (ii), the isomorphism is obtained by composing the isomorphism of Proposition 5.3 and the map induced in cohomology by the symbol map σ from $\mathcal{D}^{n-1}(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$.

To complete Theorem 6.1, we need to compute the connecting homomorphism θ .

Lemma 6.3 If $n > 0$, identifying $H(s\ell_{m+1}, C_\infty(\mathbb{R}^m, S_{q+n}^{n,q}\mathbb{R}^m))$ and $H(s\ell_{m+1}, \mathcal{D}^{n-1}(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)))$ with $(\Lambda gl(m, \mathbb{R})^*)_{g-inv}$ using Proposition 5.3 and Theorem 6.1, one has

$$\theta(\gamma) = (-1)^{t+1} n[(m+1)\delta - (2q+n+m)]\gamma, \quad \forall \gamma \in (\Lambda^t gl(m, \mathbb{R})^*)_{g-inv}.$$

Proof. To perform the computation, it is convenient to identify the spaces $\mathcal{D}^k(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ and $\oplus_{0 \leq i \leq k} C_\infty(\mathbb{R}^m, S_p^{i,q}\mathbb{R}^m)$ by representing the operator

$$D : T \rightarrow \sum_{|\alpha| \leq k} A_\alpha (D_x^\alpha T)$$

where $A_\alpha \in C_\infty(\mathbb{R}^m, Hom(\vee^p \mathbb{R}^m, \vee^q \mathbb{R}^m))$, by

$$\mathcal{T}_D : (\eta, P) \in \mathbb{R}^{m*} \times \vee^p \mathbb{R}^m \rightarrow \sum_{|\alpha| \leq k} \eta^\alpha A_\alpha(P) \in C_\infty(\mathbb{R}^m, \vee^q \mathbb{R}^m).$$

(cf. the proof of Proposition 5.3). It follows then easily that, the components of $\zeta \in \mathbb{R}^{m*}$ representing symbolically the partial derivatives of $X \in Vect(\mathbb{R}^m)$,

$$\begin{aligned} \mathcal{T}_{L_X D}(\eta, P) &= (X \cdot \mathcal{T}_D)(\eta, P) + \langle X, \eta \rangle \mathcal{T}_D(\eta, P) + \delta \langle X, \zeta \rangle \mathcal{T}_D(\eta, P) \\ &\quad - X(\zeta D_\xi) \mathcal{T}_D(\eta, P) - \mathcal{T}_D(\eta + \zeta, \langle X, \eta \rangle P + \delta \langle X, \zeta \rangle P - X(\zeta D_\xi) P) \end{aligned} \quad (7)$$

(see [9] where similar symbolical computations are used).

Now $\gamma \in (\Lambda gl(m, \mathbb{R})^*)_{g-inv}$ defines an element in $H(s\ell_{m+1}, C_\infty(\mathbb{R}^m, S_{q+n}^{m,q} \mathbb{R}^m))$, namely the class of the cocycle $c = \chi(\gamma)$

$$(X_0, \dots, X_{t-1}) \mapsto (-1)^t \gamma(DX_0, \dots, DX_{t-1})(\eta D_\xi)^n.$$

In view of the above identification, c may be considered as valued in $\mathcal{D}^n(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$. Its coboundary is valued in $\mathcal{D}^{n-1}(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ and we seek for the $\gamma' \in (\Lambda gl(m, \mathbb{R})^*)_{g-inv}$ which $[\partial c]$ corresponds to through the isomorphism (ii) of Theorem 6.1 (see Remark 6.2). To compute γ' , it suffices to evaluate $(\partial c)(A_0^*, \dots, A_{t-1}^*, \alpha^*)$, $A_i \in gl(m, \mathbb{R})$, $\alpha \in \mathbb{R}^m$, at $x = 0$. That is the same to compute the constant term of

$$(-1)^t L_{\alpha^*}(c(A_0^*, \dots, A_{t-1}^*)) \quad (8)$$

where L_{α^*} is the derivation of differential operators in the direction of α^* . Observe that

$$c(A_0^*, \dots, A_{t-1}^*) = \gamma(A_0, \dots, A_{t-1})(\eta D_\xi)^n$$

has constant coefficients so that we need only the terms of (8) involving second order derivatives of the coefficients of α^* . It follows from (7) that if $X = \alpha^*$, the terms of second order in X in $\mathcal{T}_{L_X D}(\eta, P)$ are given by

$$\begin{aligned} &-(\eta D_\eta)(\alpha D_\eta) \mathcal{T}_D(\eta, P) - (m+1)\delta(\alpha D_\eta) \mathcal{T}_D(\eta, P) \\ &+(\alpha D_\eta) \mathcal{T}_D(\eta, (\xi D_\xi)P) + \sum_i D_{\eta_i} \mathcal{T}_D(\eta, \xi_i(\alpha D_\xi)P). \end{aligned}$$

With $\mathcal{T}_D(\eta, P) = (\eta D_\xi)^n P$ and $P \in \vee^{q+n} \mathbb{R}^m$, this gives

$$-n[(m+1)\delta - (2q+n+m)](\alpha D_\xi)(\eta D_\xi)^{n-1}.$$

Hence the lemma. ■

Theorem 6.4 *If $n > 0$, the space $H(s\ell_{m+1}, \mathcal{D}^n(\mathcal{S}_\delta^{q+n}(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)))$ is isomorphic to $(\Lambda gl(m, \mathbb{R})^*)_{g-inv}^2$ if $\delta = (2q+n+m)/(m+1)$ and vanishes otherwise. If $n = 0$, it is isomorphic to $(\Lambda gl(m, \mathbb{R})^*)_{g-inv}$.*

Proof. This follows immediately from (6) and previous results. ■

We are now able to compute the cohomology of $s\ell_{m+1}$ with coefficients in $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$.

Theorem 6.5 *If $p < q$, then $H(s\ell_{m+1}, \mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))) = 0$. Otherwise, the inclusion*

$$\mathcal{D}^n(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)) \xrightarrow{i} \mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$$

induces an isomorphism in cohomology, where $n = p - q \geq 0$.

Proof. For the sake of simplicity, we denote here by $\mathcal{D}$ and $\mathcal{D}^k$ the spaces $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ and $\mathcal{D}^k(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ respectively.

Observe that a cochain on $s\ell_{m+1}$ valued in the space of differential operators necessarily takes its values in the space of operators of a certain order k . This is because $s\ell_{m+1}$ is finite dimensional.

This implies first that $H(s\ell_{m+1}, \mathcal{D}) = 0$ if $p < q$ because, in this case, the spaces $H(s\ell_{m+1}, \mathcal{D}^k)$ are all vanishing.

Second, this also implies that if $p = q + n$, $n \geq 0$,

$$i_\# : H(s\ell_{m+1}, \mathcal{D}^n) \rightarrow H(s\ell_{m+1}, \mathcal{D})$$

is an isomorphism. Indeed, if $i_\#[S] = 0$, then S is the coboundary of some $\mathcal{D}^k$ -valued cochain T . If $k \leq n$, then $[S] = 0$ because $\mathcal{D}^k \subset \mathcal{D}^n$. If $k > n$, then again $[S] = 0$, due to Theorem 6.1 (b) (iii). This proves that $i_\#$ is injective. A similar discussion shows that it is onto. ■

Remark 6.6 When $p > q$, the cohomology of $s\ell_{m+1}$ with coefficients in $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ is non vanishing if and only if $\delta = \frac{m+p+q}{m+1}$. This is why we say that $\frac{m+p+q}{m+1}$ is a *critical value* for δ . When $m = 1$, these critical values are the special values pointed out in [6] in classifying the $s\ell_2$ -module $\mathcal{D}_{\lambda\mu}^k(\mathbb{R})$. In [6], they were called “resonant” instead of critical.

For $n \geq 0$, we define

$$\tau_n : s\ell_{m+1} \rightarrow \mathcal{D}(\mathcal{S}_\delta^{*+n}(\mathbb{R}^m), \mathcal{S}_\delta^*(\mathbb{R}^m))$$

by

$$\tau_n(X)P = - \sum_{j, i_1, \dots, i_n} \partial_j X^j \partial_{i_1} \dots \partial_{i_n} D_{\xi_{i_1}} \dots D_{\xi_{i_n}} P$$

(recall that ∂_i denotes the partial derivation with respect to the i -th coordinate and that D_{ξ_j} denotes the derivation with respect to the j -th component of $\xi \in \mathbb{R}^{m*}$).

Using the notations of the proof of Lemma 6.3, one can also write

$$\tau_n(X) = - \langle X, \zeta \rangle (\eta D_\xi)^n$$

where η and ζ represent the derivatives acting on P and X respectively.

For $n > 0$, we also introduce the map

$$\gamma_n : s\ell_{m+1} \rightarrow \mathcal{D}(\mathcal{S}_\delta^{*+n}(\mathbb{R}^m), \mathcal{S}_\delta^*(\mathbb{R}^m))$$

given by

$$\gamma_n(X)P = \frac{1}{m+1} \sum_{j, i_1, \dots, i_n} \partial_{i_1} \partial_j X^j \partial_{i_2} \dots \partial_{i_n} D_{\xi_{i_1}} \dots D_{\xi_{i_n}} P.$$

It is symbolically given by

$$\gamma_n(X) = \frac{1}{m+1} \langle X, \zeta \rangle (\zeta D_\xi)(\eta D_\xi)^{n-1}.$$

Corollary 6.7 *Let $p = q + n$. The space $H^1(s\ell_{m+1}, \mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)))$ is spanned by $[\tau_0]$ if $n = 0$ and by $[\tau_n]$ and $[\gamma_n]$ if $n > 0$ and $\delta = \frac{m+p+q}{m+1}$. It vanishes otherwise.*

Proof. This easily follows from the previous results. ■

7 The space $Hom_{s\ell_{m+1}}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$

In this section, we compute the 0-th cohomology space of $s\ell_{m+1}$ with coefficients in $Hom(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$. It is the space $Hom_{s\ell_{m+1}}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ of $s\ell_{m+1}$ -equivariant linear mappings from $\mathcal{S}_\delta^p(\mathbb{R}^m)$ into $\mathcal{S}_\delta^q(\mathbb{R}^m)$.

For each $n \in \mathbb{N}$, we define

$$T_n : \mathcal{S}_\delta^{*+n}(\mathbb{R}^m) \rightarrow \mathcal{S}_\delta^*(\mathbb{R}^m)$$

by

$$T_n P = \sum_{i_1, \dots, i_n} \partial_{i_1} \dots \partial_{i_n} D_{\xi_{i_1}} \dots D_{\xi_{i_n}} P.$$

With the notations of the proof of Lemma 6.3, T_n is represented by the polynomial $(\eta D_\xi)^n$. Moreover, it follows from that proof that

$$L_X \circ T_n - T_n \circ L_X = -n[(m+1)\delta - (m+2p+n)]\gamma_n(X).$$

In particular, $T_n : \mathcal{S}_\delta^{p+n}(\mathbb{R}^m) \rightarrow \mathcal{S}_\delta^p(\mathbb{R}^m)$ is $s\ell_{m+1}$ -equivariant if and only if $n[(m+1)\delta - (m+2p+n)] = 0$.

Observe that

$$\tau_n(X) = -\langle X, \zeta \rangle T_n$$

and

$$\gamma_n(X) = \frac{1}{m+1} \langle X, \zeta \rangle (\zeta D_\xi) \circ T_{n-1}.$$

Moreover,

$$\gamma_{n+i}(X) = \gamma_n(X) \circ T_i$$

since $T_i \circ T_j = T_{i+j}$.

Lemma 7.1 *If $p \geq q$ and if $A \in Hom(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ commute with the Lie derivations in the direction of $\partial_1, \dots, \partial_m$ and of $E = x^i \partial_i$ then A is a differential operator.*

Proof. It has been shown in [8] that if $\delta = 0$, then under the assumptions of the lemma, A is a local operator. In fact, the proof of [8] works also if $\delta \neq 0$. Thus, we may assume that A is local. From the theorem of Peetre [10], it is then locally a differential operator. Since it commutes with $L_{\partial_1}, \dots, L_{\partial_m}$, its order is bounded. ■

Theorem 7.2 *If $p \geq q$ and if $A \in Hom_{s\ell_{m+1}}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ is non vanishing then either $p = q$ and A is a constant multiple of the identity or $p > q$, $\delta = \frac{m+p+q}{m+1}$ and A is a constant multiple of T_{p-q} .*

Proof. Indeed, since A is a differential operator, it is a 0-cocycle of sl_{m+1} valued in $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$. The theorem then follows immediately from Theorem 6.5. $\blacksquare$

The case $p < q$ is more difficult. We state directly the theorem. Indeed, we do not know whether a lemma similar to Lemma 7.1 holds true in this case.

Theorem 7.3 *If $p < q$ and $A \in Hom_{sl_{m+1}}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$, then $A = 0$.*

Proof. It suffices to show that A is local (and thus a differential operator, proceeding like in the proof of Lemma 7.1). Indeed, it follows from Theorem 6.5 that a 0-cocycle of sl_{m+1} with values in $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m))$ is vanishing, because $p < q$.

The proof has three parts. In the first, δ has not the critical value $\frac{m+p+q}{m+1}$ and we show directly that $A = 0$. In the second, $\delta = \frac{m+p+q}{m+1}$ and $m > 1$ while in the third, $\delta = \frac{m+p+q}{m+1}$ but $m = 1$. We set again $q = p + n$.

(i) *The non critical case.* We make use of the Casimir operator $\mathcal{C}_\delta^p$ of the sl_{m+1} -module $\mathcal{S}_\delta^p(\mathbb{R}^m)$. Recall from [1] that it is the sl_{m+1} -equivariant linear map from $\mathcal{S}_\delta^p(\mathbb{R}^m)$ into itself defined by

$$\mathcal{C}_\delta^p = \sum_i L_{X_i^*} \circ L_{Y_i^*}$$

where $\{X_i : i \leq m(m+2)\}$ is any basis of $sl(m+1, \mathbb{R})$ and $\{Y_i : i \leq m(m+2)\}$ is its dual with respect to the Killing form $\mathbf{K}$ of $sl(m+1, \mathbb{R})$ (i.e. $\mathbf{K}(X_i, Y_j) = \delta_{ij}$). By Theorem 7.2, $\mathcal{C}_\delta^p$ is a multiple c_δ^p of the identity. It is easy to compute. One gets

$$2(m+1)c_\delta^p = m(m+1)\delta^2 - (m+2p)(m+1)\delta + 2p(m+p).$$

Since A is sl_{m+1} -equivariant, one has $A \circ \mathcal{C}_\delta^p = \mathcal{C}_\delta^{p+n} \circ A$. But

$$c_\delta^{p+n} - c_\delta^p = -n \left(\delta - \frac{m+2p+n}{m+1} \right).$$

Therefore, if $\delta \neq \frac{m+p+q}{m+1}$, then $A = 0$.

(ii) *The critical case* $\delta = \frac{m+2p+n}{m+1}$, $m > 1$. This case is more delicate to handle. We need some preparations. For each $\alpha \in \mathbb{R}^{m*}$, $\hat{\alpha}$ denotes the function $x \rightarrow \alpha(x)$ on $\mathbb{R}^m$ (it is α but viewed as an element of $C_\infty(\mathbb{R}^m)$). Moreover, $I \in \mathcal{S}_0^1(\mathbb{R}^m)$ is defined by

$$I(\xi)_x = \xi(x), \quad \forall x \in \mathbb{R}^m, \quad \forall \xi \in \mathbb{R}^{m*}.$$

One has

$$L_{\alpha^*} Q = L_E(\hat{\alpha} Q) + (\delta - 1)\hat{\alpha} Q - I(\alpha D_\xi) Q, \quad \forall Q \in \mathcal{S}_\delta^*(\mathbb{R}^m).$$

Using this, the sl_{m+1} -equivariance of A and the assumption $\delta = \frac{m+2p+n}{m+1}$, one gets, after some computations,

$$(m+p-1)A(\hat{\alpha} Q)_0 = A(I(\alpha D_\xi) Q)_0, \quad \forall Q \in \mathcal{S}_\delta^p(\mathbb{R}^m),$$

the index 0 denoting the evaluation at $x = 0$. This implies easily that

$$A(\hat{\alpha} I^k Q)_0 = \frac{1}{m+p-(k+1)} A(I^{k+1}(\alpha D_\xi)Q)_0.$$

Hence, by induction on k ,

$$A(\hat{\alpha}_1 \dots \hat{\alpha}_k Q)_0 = \frac{1}{(m+p-1) \dots (m+p-k)} A(I^k(\alpha_1 D_\xi) \dots (\alpha_k D_\xi)Q)_0$$

for each $Q \in \mathcal{S}_\delta^p(\mathbb{R}^m)$ and for all $\alpha_1, \dots, \alpha_k \in \mathbb{R}^{m*}$.

We now prove that A is local. Suppose that $P \in \mathcal{S}_\delta^p(\mathbb{R}^m)$ vanishes in a neighborhood of $y \in \mathbb{R}^m$. We want to show that $A(P)_y = 0$.

Assume first that $y = 0$. There exists then $Q_{i_0 \dots i_p} \in \mathcal{S}_\delta^p(\mathbb{R}^m)$ such that

$$P = \widehat{\varepsilon^{i_0}} \dots \widehat{\varepsilon^{i_p}} Q_{i_0 \dots i_p}$$

($\varepsilon^1, \dots, \varepsilon^m$ denotes canonical basis of $\mathbb{R}^{m*} : \varepsilon^i(x) = x^i, \forall x \in \mathbb{R}^m$). Then

$$A(P)_0 = \frac{1}{(m+p-1) \dots (m-1)} A(I^{p+1} D_{\xi_{i_0}} \dots D_{\xi_{i_p}} Q_{i_0 \dots i_p})_0 = 0$$

because the $Q_{i_0 \dots i_p}$'s are homogeneous of order p in $\xi \in \mathbb{R}^{m*}$.

One reduces the case $y \neq 0$ to the case $y = 0$ just by replacing above x by $x - y$ in $\hat{\alpha}, I, E, \alpha^*$, etc.

(iii) *The critical case* $\delta = \frac{m+2p+n}{m+1}$, $m = 1$. We denote by t the canonical coordinate of $\mathbb{R}$. The map A is of the form

$$A : f \left(\frac{d}{dt} \right)^p |dt|^\delta \rightarrow A_f \left(\frac{d}{dt} \right)^{p+n} |dt|^\delta, \quad f, A_f \in C_\infty(\mathbb{R}).$$

Expressing the fact that A commutes with $L_{\frac{d}{dt}}$ and $L_{t \frac{d}{dt}}$ leads immediately to the following relations :

$$\frac{d}{dt} A_f = A_{\frac{df}{dt}}$$

and

$$t \frac{d}{dt} A_f = A_{t \frac{df}{dt}} + n A_f.$$

From this, it follows first that $A_f = 0$ if f is a polynomial. Indeed, the first relation shows that if f is a homogeneous polynomial of degree s , then A_f is a polynomial of degree $\leq s$ but the second relation shows that, in the same time, A_f is homogeneous of order $s + n > s$.

On the other hand, $T_n : \mathcal{S}_\delta^{p+n}(\mathbb{R}) \rightarrow \mathcal{S}_\delta^p(\mathbb{R})$ is $s\ell_2$ -equivariant (because δ is critical). By Theorem 7.2, it follows that $A \circ T_n$ is a constant multiple a of the identity on $\mathcal{S}_\delta^{p+n}(\mathbb{R})$. It is clear that $a = 0$ because A vanishes on polynomials. This reads

$$A_{\frac{d^n}{dt^n} f} = 0$$

for all $f \in C_\infty(\mathbb{R})$. Therefore, $A = 0$. ■

8 The splitting of the short exact sequence (1) for $M = R^m$

Our aim in this section is to obtain a necessary and sufficient condition for the short exact sequence of sl_{m+1} -modules (1) to be split (with $M = \mathbb{R}^m$).

We denote by $\varphi : \mathcal{S}_\delta(\mathbb{R}^m) \rightarrow \mathcal{D}_{\lambda\mu}(\mathbb{R}^m)$ the canonical right inverse of the symbol map. If $P \in \mathcal{S}_\delta^k(\mathbb{R}^m)$, then $\varphi(P) \in \mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m)$ is the unique homogeneous differential operator of order k such that $\sigma(\varphi(P)) = P$. As known, the coboundary E_k of the restriction of φ to $\mathcal{S}_\delta^k(\mathbb{R}^m)$ takes its values in $Hom(\mathcal{S}_\delta^k(\mathbb{R}^m), \mathcal{D}_{\lambda\mu}^{k-1}(\mathbb{R}^m))$. Moreover, its cohomology class characterizes the isomorphism class of the short exact sequence (1) [1]. In particular, (1) is split if and only if E_k is a coboundary of some element of $Hom(\mathcal{S}_\delta^k(\mathbb{R}^m), \mathcal{D}_{\lambda\mu}^{k-1}(\mathbb{R}^m))$.

Lemma 8.1 *One has $E_k = -u_k\varphi \circ \gamma_1$, where*

$$u_k = (m+1)\lambda + k - 1.$$

Proof. It is just a matter of simple computation. ■

Lemma 8.2 *As a cocycle of sl_{m+1} valued in $Hom(\mathcal{S}_\delta^k(\mathbb{R}^m), \mathcal{S}_\delta^{k-n}(\mathbb{R}^m))$, γ_n is a coboundary if and only if*

$$v_n = -n((m+1)\delta - (m+2k-n))$$

is non vanishing.

Proof. By Corollary 6.7, it suffices to show that if $\gamma_n = \partial T$, then $T \in Hom(\mathcal{S}_\delta^k(\mathbb{R}^m), \mathcal{S}_\delta^{k-n}(\mathbb{R}^m))$ is a differential operator. Since $\gamma_n(\partial_i) = 0$ and $\gamma_n(E) = 0$, this follows immediately from Lemma 7.1. ■

Remark 8.3 It is clear that if $v_n \neq 0$, then γ_n is the coboundary of $\frac{1}{v_n} T_n$.

In the sequel, for $\mathcal{A} = \sum_i \mathcal{A}_i t^i \in \mathbb{R}[t]$, we denote by $\mathcal{A}(\eta\mathcal{D}_\xi)$ the differential operator

$$\sum_i \mathcal{A}_i T_i : \mathcal{S}_\delta^k(\mathbb{R}^m) \rightarrow \bigoplus_{o \leq k} \mathcal{S}_\delta^{k-i}(\mathbb{R}^m).$$

Proposition 8.4 *If $\gamma_1, \dots, \gamma_{n-1}$ are coboundaries, then*

$$E_k = -\frac{u_k \dots u_{k-n+1}}{v_1 \dots v_{n-1}} \varphi \circ \gamma_n + \partial\varphi \circ \mathcal{A}_n(\eta\mathcal{D}_\xi) \quad (9)$$

for some polynomial $\mathcal{A}_n$ of degree $n-1$ such that $\mathcal{A}_n(0) = 0$.

Proof. We proceed by induction on n . By Lemma 8.1, $E_k = -u_k\varphi \circ \gamma_1$. Assume that (9) holds true and that γ_n is a coboundary. Then, setting

$$a = -\frac{u_k \dots u_{k-n+1}}{v_1 \dots v_{n-1}}$$

for simplicity, one successively gets

$$\begin{aligned}
& (E_k - \partial\varphi \circ \mathcal{A}_n(\eta D_\xi))(X) \\
&= \frac{a}{v_n} (\varphi \circ L_X \circ T_n - \varphi \circ T_n \circ L_X) \\
&= \frac{a}{v_n} (\varphi \circ L_X \circ T_n - L_X \circ \varphi \circ T_n) + \frac{a}{v_n} (L_X \circ \varphi \circ T_n - \varphi \circ T_n \circ L_X) \\
&= -\frac{a}{v_n} E_{k-n}(X) \circ T_n + \frac{a}{v_n} (\partial(\varphi \circ T_n))(X) \\
&= \frac{au_{k-n}}{v_n} \varphi \circ \gamma_1(X) \circ T_n + \frac{a}{v_n} (\partial(\varphi \circ T_n))(X) \\
&= \frac{au_{k-n}}{v_n} \varphi \circ \gamma_{n+1}(X) + \frac{a}{v_n} (\partial(\varphi \circ T_n))(X)
\end{aligned}$$

because $\gamma_1 \circ T_n = \gamma_{n+1}$. ■

Theorem 8.5 *The short exact sequence of sl_{m+1} -modules*

$$0 \rightarrow \mathcal{D}_{\lambda\mu}^{k-1}(\mathbb{R}^m) \xrightarrow{i} \mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m) \xrightarrow{\sigma} \mathcal{S}_\delta^k \rightarrow 0 \quad (k \geq 1)$$

is split if and only if

$$\delta \notin \left\{ \frac{m+k}{m+1}, \frac{m+k+1}{m+1}, \dots, \frac{m+2k-1}{m+1} \right\} \quad (10)$$

or

$$\delta = \frac{m+2k-n}{m+1} \quad \text{and} \quad \lambda = \frac{i-k}{m+1} \quad (11)$$

for some $n \in \{1, \dots, k\}$ and some $i \in \{1, \dots, n\}$.

Proof. Let n denote the least integer such that γ_n is not a coboundary, as a cocycle valued in $Hom(\mathcal{S}_\delta^k(\mathbb{R}^m), \mathcal{S}_\delta^{k-n}(\mathbb{R}^m))$ (we set $n = k+1$ if $\gamma_1, \dots, \gamma_k$ are coboundaries).

It follows from Lemma 8.2 and Proposition 8.4 that E_k is a coboundary if (10), or (11), holds true.

Conversely, assume that E_k is a coboundary and that $n \leq k$. Then, $\delta = \frac{m+2k-n}{m+1}$. Moreover, it follows from (8) that $u_k \dots u_{k-n+1} \gamma_n$ is a coboundary. Since γ_n is not a coboundary, $u_i = 0$ for some $i \in \{k-n+1, \dots, k\}$. Hence the result. ■

Corollary 8.6 *If $(m+1)\delta - m \notin \mathbb{N}_0$, then there exists a unique sl_{m+1} -equivariant linear bijection $\sigma^{\lambda\mu} : \mathcal{D}_{\lambda\mu}(\mathbb{R}^m) \rightarrow \mathcal{S}_\delta(\mathbb{R}^m)$ such that, for each $A \in \mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m)$, the term of highest order of $\sigma^{\lambda\mu}(A)$ is the symbol $\sigma(A)$ of A .*

Proof. The existence of $\sigma^{\lambda\mu}$ follows immediately from Theorem 8.5. For the uniqueness, assume that σ' has the same properties than $\sigma^{\lambda\mu}$. Then, for each k , the restriction of $S = \sigma' \circ (\sigma^{\lambda\mu})^{-1}$ to $\mathcal{S}_\delta^k(\mathbb{R}^m)$ is of the form

$$P \in \mathcal{S}_\delta^k(\mathbb{R}^m) \rightarrow (P, S_1(P), \dots, S_k(P)) \in \bigoplus_{0 \leq \ell \leq k} \mathcal{S}_\delta^\ell(\mathbb{R}^m)$$

for some sl_{m+1} -equivariant $S_i \in Hom(\mathcal{S}_\delta^k(\mathbb{R}^m), \mathcal{S}_\delta^{k-i}(\mathbb{R}^m))$, $i = 1, \dots, k$. By Theorem 7.2, $S_i = 0$ for $i = 1, \dots, k$. ■

Remark 8.7 The existence and uniqueness of $\sigma^{\lambda\mu}$ was shown in [8] for $\lambda = \mu$ and $m > 1$. It has been obtained in [6, 2] for $m = 1$ and non critical values of δ . For critical δ , $\mathcal{D}_{\lambda\mu}(\mathbb{R})$ has been described in [6]. We will not discuss here the case of the critical values of δ in higher dimension. The next result is also a generalization of a useful result of [6, 8].

Proposition 8.8 *If $(m+1)\delta - m \notin \mathbb{N}_0$, then $T : \mathcal{D}_{\lambda\mu}(\mathbb{R}^m) \rightarrow \mathcal{D}_{\lambda'\mu'}(\mathbb{R}^m)$ is $s\ell_{m+1}$ -equivariant if and only if there exists constant $a_k \in \mathbb{R}$ such that, for each $k \in \mathbb{N}$,*

$$\sigma^{\lambda'\mu'} \circ T \circ (\sigma^{\lambda\mu})^{-1} = a_k \text{id}$$

on $\mathcal{S}_\delta^k(\mathbb{R}^m)$.

Proof. This follows immediately from Theorem 7.2 and Theorem 7.3. ■

9 Cohomology of $s\ell_{m+1}$ valued in $\mathcal{D}_{\lambda\mu}(\mathbb{R}^m)$

The computation of $H(s\ell_{m+1}, \mathcal{D}_{\lambda\mu}(\mathbb{R}^m))$ is quite similar to that of $H(s\ell_{m+1}, \mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}^m), \mathcal{S}_\delta^q(\mathbb{R}^m)))$. It uses the short exact sequence (1) and Proposition 5.1 to get first the spaces $H(s\ell_{m+1}, \mathcal{D}_{\lambda\mu}^k(\mathbb{R}^m))$, $k \in \mathbb{N}$. In most of the cases, the cohomology of the module $C_\infty(\mathbb{R}^m, \mathcal{S}_\delta^k \mathbb{R}^m)$ vanishes. It is however necessary to compute the connecting homomorphism associated to the sequence (1) for some values of δ and k . This leads also to a sort of critical values for λ and μ . In view of Proposition 5.1, they are not the same when $m > 1$ as when $m = 1$.

We only summarize the results, leaving the reader to supply the proofs.

A. *The case $m > 1$*

Theorem 9.1 $\{m > 1\}$ *If $\delta \notin \{0, 1\}$ or if $\delta = 1$ and $\lambda \neq 0$, then $H(s\ell_{m+1}, \mathcal{D}_{\lambda\mu}(\mathbb{R}^m)) = 0$.*

Theorem 9.2 $\{m > 1\}$ *The space $H(s\ell_{m+1}, \mathcal{D}_{\lambda\lambda}^0(\mathbb{R}^m))$ is isomorphic to $(\Lambda gl(m, \mathbb{R})^*)_{g\text{-inv}}$. The inclusion of $\mathcal{D}_{\lambda\lambda}^0(\mathbb{R}^m)$ into $\mathcal{D}_{\lambda\lambda}(\mathbb{R}^m)$ induces an isomorphism in cohomology.*

Theorem 9.3 $\{m > 1\}$ *One has a short exact sequence*

$$0 \rightarrow H(s\ell_{m+1}, \mathcal{D}_{01}^0(\mathbb{R}^m)) \xrightarrow{i_\#} H(s\ell_{m+1}, \mathcal{D}_{01}^1(\mathbb{R}^m)) \xrightarrow{\sigma_\#} H(s\ell_{m+1}, C_\infty(\mathbb{R}^m, S_1^1 \mathbb{R}^m)) \rightarrow 0$$

where the kernel and the quotient are isomorphic to $(\Lambda gl(m, \mathbb{R})^*)_{g\text{-inv}}$. The inclusion of $\mathcal{D}_{01}^1(\mathbb{R}^m)$ into $\mathcal{D}_{01}(\mathbb{R}^m)$ induces an isomorphism in cohomology.

Remark 9.4 If $\lambda \neq 0$, then $H(s\ell_{m+1}, \mathcal{D}_{\lambda, \lambda+1}^0(\mathbb{R}^m))$ is isomorphic to $(\Lambda gl(m, \mathbb{R})^*)_{g\text{-inv}}$ while $H(s\ell_{m+1}, \mathcal{D}_{\lambda, \lambda+1}^1(\mathbb{R}^m)) = 0$.

B. *The case $m = 1$*

In this case, the “critical” values are given by $(\lambda, \mu) = \left(\frac{1-n}{2}, \frac{1+n}{2}\right)$, $n \in \mathbb{N}_0$. Moreover, the description of the cohomology is very close to that of the module $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}), \mathcal{S}_\delta^q(\mathbb{R}))$. This follows from the fact that this module is isomorphic to $\mathcal{D}_{\delta-p, \delta-q}(\mathbb{R})$ since $\mathcal{S}_\delta^p(\mathbb{R})$ is isomorphic to the space of $(\delta-p)$ -densities. In particular, the critical values introduced in Section 6 occur when $p - q$ is a positive integer n . They are then given by $\delta = (p + q + 1)/2$. In this case,

$$\delta - p = \frac{1-n}{2} \quad \text{and} \quad \delta - q = \frac{1+n}{2}.$$

Note that these critical values were also obtained in [6], but in a completely different way.

Theorem 9.5 *If $\delta \notin \mathbb{N}$ or if $\delta \in \mathbb{N}_0$ and $\lambda \neq \frac{1-\delta}{2}$, then $H(\mathfrak{sl}_2, \mathcal{D}_{\lambda\mu}(\mathbb{R})) = 0$.*

Theorem 9.6 *The inclusion of $\mathcal{D}_{\lambda\lambda}^0(\mathbb{R})$ into $\mathcal{D}_{\lambda\lambda}(\mathbb{R})$ induces an isomorphism in cohomology. In particular*

$$H^i(\mathfrak{sl}_2, \mathcal{D}_{\lambda\lambda}(\mathbb{R})) = \begin{cases} \mathbb{R} & \text{if } i = 0, 1 \\ 0 & \text{if } i = 2, 3 \end{cases}$$

Theorem 9.7 *If $\lambda = \frac{1-n}{2}$ and $\mu = \frac{1+n}{2}$ for some $n \in \mathbb{N}_0$, then for each u , one has a short exact sequence*

$$0 \rightarrow H^u(\mathfrak{sl}_2, \mathcal{D}_{\lambda\mu}^{n-1}(\mathbb{R})) \xrightarrow{i_\#} H^u(\mathfrak{sl}_2, \mathcal{D}_{\lambda\mu}^n(\mathbb{R})) \xrightarrow{\sigma_\#} H^u(\mathfrak{sl}_2, C_\infty(\mathbb{R}, S_n^*\mathbb{R})) \rightarrow 0$$

where the kernel (resp. the quotient) is isomorphic to $\mathbb{R}$ for $u = 1, 2$ (resp. $0, 1$) and is vanishing otherwise. Moreover, the inclusion of $\mathcal{D}_{\lambda\mu}^n(\mathbb{R})$ into $\mathcal{D}_{\lambda\mu}(\mathbb{R})$ induces an isomorphism in cohomology.

Remark 9.8 1) One sees in particular that $H^3(\mathfrak{sl}_2, \mathcal{D}_{\lambda\mu}(\mathbb{R})) = 0$ for all $\lambda, \mu \in \mathbb{R}$.

2) For $(\lambda, \mu) = \left(\frac{1-n}{2}, \frac{1+n}{2}\right)$, $n \in \mathbb{N}_0$, $H^1(\mathfrak{sl}_2, \mathcal{D}_{\lambda\mu}(\mathbb{R}))$ is spanned by

$$\tau'_n \left(f \frac{d}{dt} \right) : g |dt|^\lambda \rightarrow \frac{df}{dt} \frac{d^n g}{dt^n} |dt|^\mu$$

and

$$\gamma'_n \left(f \frac{d}{dt} \right) : g |dt|^\lambda \rightarrow \frac{d^2 f}{dt^2} \frac{d^{n-1} g}{dt^{n-1}} |dt|^\mu.$$

This follows from Theorem 9.7. The γ'_n 's are exactly the cocycles used in [6] in the study of the $\mathfrak{sl}_2$ -structure of $\mathcal{D}_{\lambda\mu}(\mathbb{R})$.

3) If $p = q + n$ and $\delta = \frac{p+q+n}{2}$, through the isomorphism between $\mathcal{D}(\mathcal{S}_\delta^p(\mathbb{R}), \mathcal{S}_\delta^q(\mathbb{R}))$ and $\mathcal{D}_{\frac{1-n}{2}, \frac{1+n}{2}}(\mathbb{R})$, τ_n and γ_n become respectively τ'_n and γ'_n .

Acknowledgements

We would like to thank very much M. De Wilde, P. Mathonet and V. Ovsienko for helpful discussions.

References

- [1] Y. Chow. *General theory of Lie algebras*, Vol. 1 and Vol. 2. Gordon and Breach. New York, London, Paris (1978).
- [2] P. Cohen, Y. Manin, D. Zagier. Automorphic pseudodifferential operators. Preprint MPI (1995).
- [3] C. Duval, V. Ovsienko. Space of second order linear differential operators as a module over the Lie algebra of vector fields. To appear in *Advances in Math.*
- [4] D.B. Fuks. *Cohomology of Infinite Dimensional Lie Algebras*. Contemporary Soviet Mathematics. Consultant Bureau. New York and London (1986).
- [5] H. Gargoubi, V. Ovsienko. Space of linear differential operators on the real line as a module over the Lie algebra of vector fields. *Int. Res. Math. Notes*, 5 (1996), pp. 235–251.
- [6] H. Gargoubi. Modules des opérateurs différentiels sur la droite : géométrie projective et cohomologie de Gelfand-Fuks. Thèse de doctorat. Université de Marseille Provence I (1997).
- [7] W. Greub, S. Halperin, R. Vanstone. *Connections, curvature and cohomology*. Vol. I. Pure and Applied Mathematics 47. Academic Press. New York and London (1972).
- [8] P. Lecomte, V. Ovsienko. Projectively invariant symbol map and cohomology of vector fields Lie algebras intervening in quantization. Preprint CPT-CNRS Luminy (1996).
- [9] P. Lecomte, P. Mathonet, E. Tousset. Comparison of some modules of the Lie algebra of vector fields. *Indag. Mathem., N.S.*, 7(4) (1996), pp. 461–471.
- [10] J. Peetre. Une caractérisation des opérateurs différentiels. *Math. Scand.* **7** (1959), pp. 211–218 and **8** (1960), pp. 116–120.
- [11] H. Weyl. *The classical groups, their invariants and representations*. Princeton Math. Series. Princeton (1946).

P. LECOMTE
Institut de Mathématique
Université de Liège
Grande Traverse 12, B37
B-4000 Liège (Belgium)

e-mail : plecomte@ulg.ac.be